Ph:+91 844-844-0102

1.1 Linear Algebra: Why And How

Why?

- Huge Applications in ML/AI
- A lot of applications in Physics (Velocity, accelerations, force)
- In all fields of Engineering (Mechanical, Civil, electrical, etc).
- For example in computer science
  1. Computer Graphics → screen, games, edit an image.
  2. Linear System of Equations. (solutions to linear system of Equns)
  3. Data Compression.
  4. Recommender Systems.
- Also have a lot of applications in Physics, Chemistry, Biology.

How

- Usually L.A. is taught in Numeric/Mathematic way.
- Geometric/Intuitive way helps us visualize/ picturize the concept.
- The geometric approach provides us a much deeper understanding and is fun to learn.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Book :- Linear Algebra and Its Applications - Gilbert Strang.

Video lectures :- MIT lectures on Youtube by Gilbert Strang.

3 blue 1 brown :- Essence of Linear Algebra, for visualizations.
↓
Youtube Channel

Ph: +91 844-844-0102

1.2. How To Learn Mathematics

1. Abstract / Mathematical way of learning → Mathematics.
2. Applied way of learning → Scientist Engineers.

eg Prime Numbers → Applications in Cryptography :- Logins on websites securely //
↓
Mathematicians.

Applied Mathematicians / Scientists

- Mathematics - consist of equations but it has a lot of meaning behind it like poetry, but we should understand the concept behind it.

- Geometric / Diagrammatic way → art/painting

- Our focus should be on both the equations and the diagrams and how to relate them both.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

1.3 Vectors - I

- Vectors
  - Encountered in Physics

Physics Interpretation:
- A vector has length/magnitude & direction (vel, acc, force).

- In terms of a computer science - a vector can be represented by arrays,

$$\begin{bmatrix} 160 \\ 55 \\ 7 \end{bmatrix}$$
- 160 — height
- 55 — Weight
- 7 — Shoe Size.

- In terms of Mathematicians :-
  - A vector is considered as an abstract concept where we have two operations 1. Vector addition 2. Scalar Multiplication

Visualisation



Vector can be represented as array of dot/point, ordered list based on the application.

$$\|\vec{a}\| = \sqrt{a_1^2 + a_2^2} \qquad \vec{a} = \begin{bmatrix} a_1 \\ a_2 \end{bmatrix}$$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

→ The concepts of 2D, 3D can be extended to higher dimension space like 4D, 5D ... 10D or even 1000D.

Ph: +91 844-844-0102

- Addition of Vectors

$$V_1 = \begin{bmatrix} 2 \\ 3 \end{bmatrix} \quad V_2 = \begin{bmatrix} 4 \\ 5 \end{bmatrix} \rightarrow = \begin{bmatrix} 6 \\ 8 \end{bmatrix}$$

$$\begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} + \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix} = \begin{bmatrix} a_1 + b_1 \\ a_2 + b_2 \\ \vdots \\ a_n + b_n \end{bmatrix}$$ n Dimensional.

Geometric Interpretation.



$$\vec{c} = \vec{a} + \vec{b}$$

$$\vec{c} = \begin{bmatrix} a_1 + b_1 \\ a_2 + b_2 \end{bmatrix}$$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

1-D addition

$2+3=5$



$\vec{a} + \vec{b}$

Scalar Multiplication of a vector or Multiplication of a vector By a Scalar

(More appropriate to use)

$$2 \times \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 4 \\ 6 \end{bmatrix}.$$

$2 \times \vec{a} = \vec{b}$.



→ The vector b has become streached

→ If the scalar is < 0 then direction is reversed.

- Mathematicians say that any operations i.e. vector operations can be written in terms of vector multiplications with a scalar and vector vector addition.

Gate Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

1.4 Vectors - 2

Unit Vectors



- $\hat{i}$ is unit vector in x direction
- $\hat{j}$ is unit vector in y direction
- $\|\hat{i}\| = 1 \;.\; \|\hat{j}\| = 1$ $\quad \hat{i} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ $\hat{j} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ $\quad \vec{a} = 2\hat{i} + 3\hat{j}$

Standard basis unit vector

$$\vec{a} = \begin{bmatrix} 2 \\ 3 \end{bmatrix} = 2\hat{i} + 3\hat{j}$$

$$= 2\begin{bmatrix} 1 \\ 0 \end{bmatrix} + 3\begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$$

- Any pair of vectors can form a basis vectors except for 2 cases.

1) $\vec{b} = \vec{a} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$

2) $\vec{b} = \text{Constant } \vec{a}$. (In this case the span of the 2 vectors is a st line).

Span :- The span is the complete space which can be represented by linear combinations of the vector or pair of vectors.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

Span of 2 vectors in 3D space :- It is always a plane / sheet.

Span of 3 vectors in 3D space :- is a 3D space.

Linear dependence :-

- If a vector $\bar{a}$ can be expressed as sum of two other vectors multiplied by some scalar then $\bar{a}$ is said to be linearly dependent on the other vectors. For example $\vec{a} = x\vec{b} + y\vec{c}$ $\vec{a}$ is linearly dependent on $\vec{b}$ and $\vec{c}$.

- If a and b are two linearly dependent vectors in the 2D space, then they cannot span the complete 2D space, instead they can span the complete 2D space.

→ If $\bar{a}$ and $\bar{b}$ span a d-dimention space (where $d < n$) : n is the dimention of the vectors a and b then if another vector c is added and the space spanned by a and b does not increase in dimention.
then c is linearly dependent on a and b.

→ Basis vectors are a set of linearly independent vectors which span the complete space in all dimentions.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

1.5 Matrices: Linear Transformations / Functions

Transformation :- Changing vector from one form to another.



- A transformation could also represent the complete space i.e. the comple space and all the lines / vectors in the space may be transformed.

Linear Transformation

1. All the lines must remain lines (not curved)
2. Origin should remain the same.

(or)

Grid lines should remain parallel and evenly spaced.

example application :- Computer graphics. :- Rotation of given image.



Rotation is a linear transformation

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

A shear is also another type of a linear transformation



i/p

o/p parallelogram.

— A linear transformation can be expressed in the form of Matrix multiplication.



This can be expressed in the form of matrix multiplication as well.

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

Transformation matrix

Final position of $\hat{i}$ after transformation

Final position of $\hat{j}$ after transformation.

→ We can get $\vec{a_1}$ by.

$$= A \cdot \vec{a}$$

$$= \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \end{bmatrix}$$

$$= a_1 \begin{bmatrix} 0 \\ 1 \end{bmatrix} + a_2 \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$

$$= \begin{bmatrix} -a_2 \\ a_1 \end{bmatrix}$$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

Example of shear linear transformation.



$$\hat{i} \rightarrow \hat{i}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}.$$

$$\hat{j} \rightarrow \hat{j}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

Matrix of transformation can be written as $\cdot M = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$

Any vector can be obtained $\bar{a}_1$ on transformation will change to.

$$\vec{a}_2 = M \cdot \vec{a}_1$$

Order of a Matrix : A matrix if consists of n no of rows and m number of columns then it is said to be of order $m \times n$.

- Even if a matrix is a non square matrix it represents a linear transformation.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

PH: +91 844-844-0102

1.6. Matrix - Matrix Multiplication and Addition

Matrix Vector Multiplication

- Linear transformation in any dimention can be expressed in form of Matrix vector multiplication $A \cdot V = U$, in 2D, 3D, n-D.
- Also if A is a non square matrix we may represent a tranform for eg. $A_{3\times2}$ will transform a 3D vector to 2D space in form of a 2D vector.

$$A_{3\times\boxed{2}}\ V_{\boxed{2}\times1} = U_{3\times1}$$

Matrix Matrix Multiplication

- Similar to composition of functions.
- Composition of matrices like performing 2 linear transforms

$$(A(B \cdot V)) = AU = W.$$

example $\begin{bmatrix} 0 & 2 \\ 1 & 0 \end{bmatrix}\begin{bmatrix} 1 & -2 \\ 1 & 0 \end{bmatrix}\begin{bmatrix} x \\ y \end{bmatrix} = ABV = CV$

$C = AB$.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

$$\begin{bmatrix} 0 & 2 \\ 1 & 0 \end{bmatrix}\begin{bmatrix} 1 & -2 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 0\times1 + 2\times1 & 0+0 \\ 1 & -2 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 0 \\ 1 & -2 \end{bmatrix}$$

Matrix multiplication is a composition of linear transforms.

- For matrix multiplication the order of the adjacent matrix should match as follows $A_{m\times n}\ B_{n\times k} = C_{m\times k}$.

Matrix-Matrix Addition

- For addition to be done the order of the two matrices should be identical.
- Sum is calculated by adding corresponding elements.
- It can be thought as computing the sum of transformations

$M_1 \rightarrow$ Transformation-1

$M_2 \rightarrow$ Transformation-2

$(M_1 + M_2)V =$ Sum of transformations M_1 and M_2 on V.

Gate Applied course.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

Properties

1. Commutativity → $M_1 + M_2 = M_2 + M_1$

   $M_1 M_2 \neq M_2 \cdot M_1$

2. Associativity. $A + (B + C) = (A + B) + C$.

   $A \cdot (BC) = (AB) \cdot C$.

3. Distributive :. $A(B+C) = AB + AC$

   $(A+B)C = AC + BC$

   $n(AB) = (nA)x = A(Bn)$

1.7 Transpose And Dot Product

- Transpose : Obtained by interchanging the rows and columns of a matrix.

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \qquad A^T = \begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$$

Properties

1. $(A^T)^T = A$

2. $(A+B)^T = A^T + B^T$.

Gatecse@appliedcourse.com

Gate.Appliedcourse.com

Ph: +91 844-844-0102

3. $(AB)^T = B^T A^T$

4. $(cA)^T = c \cdot (A^T)$

Dot Product

$a \cdot b = a^T b$

$= \begin{bmatrix} a_1 & a_2 \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$



$a \cdot b = a_1 b_1 + a_2 b_2$

- Dot product of two matrices is column wise multiplication of corresponding elements

- Geometrically $a \cdot b$ is the length of vector a × length of vector b and cosine of angle in between both of them.

$a_1 b_1 + a_2 b_2 = \|a\| \|b\| \cos\theta$



- Another way can be is the projection of a vector in the direction of the other vector.

$a \cdot b = \|a\| \|b\| \cos\theta$

$= \|b\| \text{proj}_b a$



Gatecse@appliedcourse.com

Gate.Appliedcourse.com

Ph: +91 844-844-0102

1.8 Types Of Matrices

(1) Row Matrix / Column Matrix :-

Row Matrix having only on row, whereas a column matrix is a matrix having only one colum matrix

(2) Square Matrix :- If the no of rows and columns are equal then the matrix is known as a square matrix m=n.

(3) Symmetric Matrix :- If $A = A^T$ and A is a square matrix,

$A_{ij} = A_{ji}$

$$\begin{bmatrix} 1 & 7 & 3 \\ 7 & 4 & 5 \\ 3 & 5 & 6 \end{bmatrix}_{3\times 3}$$

1, 4, 6 :- constitute the diagonal of the matrix.

(4) Skew Symmetric Matrix :- A Square Matrix $A^T = -A$.

(5) Diagonal Matrix

- Square diagonal Matrix If all elements on diagonal Elements are non zero and remaing all Elents are zero.

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$$

Gate Applied course.com

Gatecse@appliedcourse.com

Ph: +91 844 844 0102

Rectangular Diagonal Elements :- A rectangular matrix where every non diagonal element is zero and all diagonal elements are non zero.

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \end{bmatrix}$$

(6) Identity Matrix $A_{ij} = 1$ if $i = j$
$= 0$ otherwise
and A is a square matrix

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I_{3\times 3} = I_3.$$

$IA = A$

(7) Scalar Matrix :- $A_{ij} = \lambda$ if $i = j$
$= 0$ $i \neq j$
A is a square matrix

$$\begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix}$$

(8) Zero Matrix :. All elements are zero's.

$$Z = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$A \cdot Z = Z$ (if Multiplication is valid).

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

9. Upper Triangular Matrix

- If all the elements which are not in the upper triangular matrix are zero then it is said to be an upper triangular matrix.

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix}$$

10. Lower Triangular Matrix

- If all the elements which are not in the lower triangle of the matrix are zero then it is said to be a lower triangular matrix

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 3 & 0 \\ 4 & 5 & 6 \end{bmatrix}$$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

1.9 Determinant of a Matrix

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$\det(A) = |A|$$

$$|A| = (ad - bc)$$

If A represents a linear transformation then $|A|$ represents the area of the parallelogram formed by the transformed vectors for $\hat{i}$ and $\hat{j}$.

→ After the linear transformation

$\hat{i} \xrightarrow{A} \hat{i}_1 \begin{bmatrix} a \\ c \end{bmatrix}$

$\hat{j} \xrightarrow{B} \hat{j}_1 \begin{bmatrix} b \\ d \end{bmatrix}$



$\hat{i} \xrightarrow{A} \hat{i}_1$

$\hat{j} \xrightarrow{A} \hat{j}_1$

→ Another way to think about $|A|$. Scale by which the area of any figure changes due to the linear transformation represented by A.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

example.

$$A = \begin{bmatrix} 4 & 2 \\ 2 & 1 \end{bmatrix}$$

$\hat{i}$ lands $\quad \hat{j}$ lands.



$\hat{i}_1 = 2\hat{j}_1$ They are linearly dependent.

$|A| = 0$. Also the area of the Parallelogram $= 0$.

ex 2

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \qquad |A| = 4 - 6 = -2$$

- If the sign is $-ve$ after the transformation the orientation of $\hat{i}$ and $\hat{j}$ gets changed.



Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

- Same concept if extended to 3D

$\Delta_{3\times3}$ represents the volume of the parallelopiped

A still corresponds to a linear transformation.

$$|A| = \begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} = a(ei - hf) - b(di - gf) + c(dh - eg).$$

$$= a\begin{vmatrix} e & f \\ h & i \end{vmatrix} - b\begin{vmatrix} d & f \\ g & i \end{vmatrix} + c\begin{vmatrix} d & e \\ g & h \end{vmatrix}.$$

Minors of Δ.

SARRUS RULE



$= aei + bfg + cdh - ceg - bdi - afh.$

2 cols repeated

Bold lines + sign.
dotted lines − sign.

Gatecse@appliedcourse.com

Gate.Appliedcourse.com

Ph: +91 844-844-0102

Properties of determinants.

① $|I_n| = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = 1$ $\quad I \cdot V = V$.

② $|A^T| = |A|$

③ $|A^{-1}| = \frac{1}{|A|} = |A|^{-1}$

④ $|AB| = |A||B|$.

⑤ $|cA| = c^n |A|$ — A is a $n \times n$ matrix.

⑥ Triangular Matrix A_T.

$|A_T|$ = product of the diagonal elements.

Reference links

1. https://en.wikipedia.org/wiki/Determinant
2. 3 Blue 1 Brown channel Videos.
   1. https://youtu.be/Ip3X9LOh2dk?t=523
   2. " ?t=142
   3. " t=257

APPLIED COURSE
Gate.Appliedcourse.com
Gatecse@appliedcourse.com

Ph: +91 844-844-0102

1.10 Minor, Cofactor, Adjoint, Inverse of a Matrix.

Cofactor..

$$A = \begin{bmatrix} 1 & 4 & 7 \\ 3 & 0 & 5 \\ -1 & 9 & 11 \end{bmatrix}$$

$$C = \text{cofactor}(A) = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}$$

Minor M_{11}

$$C_{11} = (-1)^{1+1} \begin{vmatrix} 0 & 5 \\ 9 & 11 \end{vmatrix}$$

$$C_{23} = (-1)^{2+3} \begin{vmatrix} 1 & 4 \\ -1 & 9 \end{vmatrix}$$

$$C_{ij} = (-1)^{i+j} M_{ij}$$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

Adjoint / Adjugate / Adjunct

1. $\text{Adj}(A) = C^T = [\text{(cofactor}(A)]^T$

2. $A \cdot \text{Adj}(A) = |A| I$.

$$A \cdot \frac{\text{Adj}(A)}{|A|} = I$$

$$\therefore A^{-1} = \frac{\text{Adj}(A)}{|A|}$$

If $|A| = 0$ inverse of A is not defined, the matrix is non invertable

→ One of the application of inverse of a matrix is to find the solution of system of linear equations.

→ Geometric intution of inverse of a matrix :-

If vector v on linear tranformation we get $\bar{v}$, the inverse tranformation helps us or is that tranformation that will help us get back

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

vector v from vector u.



→ Geometric Interpretation of Adj(A).

$$Adj(A) = |A| A^{-1}$$

↙ ↓ inverse transform of A

Scaling transformation of A.

= It is scaled transformation of the inverse transform of A, the scale is same as that of scale of A.

Algebraic Properties & Geometric Properties

① $|A^{-1}| = \frac{1}{|A|}$ or $|A|^{-1}$

② $A^{-1} = \frac{adj(A)}{|A|}$.

$\Rightarrow |A| A^{-1} = adj(A)$

~~$|A| Adj(A)$~~ $\Rightarrow \left| |A| A^{-1} \right| = |Adj(A)|$

Gate.Appliedcourse.com
Gatecse@appliedcourse.com

Ph: +91 844-844-0102

$$|A|^n |A|^{-1} = |A|^n \frac{1}{|A|} = |A|^{(n-1)}$$

$$\therefore \underline{|Adj(A)| = |A|^{n-1}}$$

(3) Singular Matrix

$|A| = 0$ A^{-1} does not exist

Ilgm area = 0.

(4) $|Adj(Adj(A))| = ?$

$|adj(A)| = |A|^{n-1}$

If $Adj(B) = B$

$|Adj(B)| = |B|^{n-1}$

$|B| = |Adj(A)|$

$= |A|^{n-1}$

$\therefore |Adj(B)| = |Adj(Adj(A))| = (|A|^{n-1})^{n-1} \Rightarrow |A|^{(n-1)^2}$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

Similarly $|Adj.(Adj.(Adj(A)))| = |A|^{(n-1)^3}$. ⑤

⑤ $|A \cdot B| = |A| \cdot |B|$.

$|A^K| = |A|^K$.

⑥.

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \qquad Adj(A) = \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

$$A^{-1} = \frac{1}{(ad-bc)} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

⑦ $Adj(0) = 0$.

⑧ $adj(I) = I$

⑨ $adj(cA) = c^{(n-1)}\, adj(A)$.

⑩ $adj(A^T) = (adj(A))^T$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

(11) $\text{adj}(A^{-1}) = (\text{adj}(A))^{-1}$

(12) $\text{Adj}(AB) = \text{Adj}(B) \cdot \text{Adj}(A)$.

(13) $(AB)^{-1} = B^{-1}A^{-1}$

(14) $\text{Adj}(A^K) = (\text{Adj}(A))^K$

(15) $(A^T)^{-1} = (A^{-1})^T$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 8413844102

1.11 Solved Problems Elementry Row and Column Operations

(1) Let X be a square matrix. Consider the following two statements on X.

I. X is invertable

II. Determenant of X is non-zero.

Which of the following is TRUE?

(A) I implies II; II does not imply I

(B) II implies I; I does not imply II

(C) I does not imply II; II does not imply I.

(D) I and II are equalent statements.

If A^{-1} exists it means $|A| \neq 0$ as $A^{-1} = \frac{Adj(A)}{|A|}$

∴ Both the statements are equalent.

(2) Perform the following operations on the below matrix

$$\begin{bmatrix} 3 & 4 & 45 \\ 7 & 9 & 105 \\ 13 & 21 & 195 \end{bmatrix}$$

Gate Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91-844-844-0102

i. Add the third row to the second row.

ii. Subtract the third column from the first column.

The determinat of the resultant matrix is _____.

Elementary Row & Column Operations

① $R_i \leftrightarrow R_j$ (or) $C_i \leftrightarrow C_j$ - determinat will chage its sign.

② $R_i \leftarrow cR_i$ (or) $C_j \leftarrow cC_j$ → Resultant determinat is multiplied by c

③ $R_i \leftarrow R_i + cR_j$ or $C_i \leftarrow C_i + cC_j$ → Resultant determinat will not chage.

Coming back to the problem.

i. $R_2 \leftarrow R_2 + R_3$

ii. $C_1 \leftarrow C_1 - C_3$

By these operations the determinat of the matrix remains unchaged.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

If we observe $C_3 = 15 \times C_1$.

$\therefore \quad C_3 \leftarrow C_3 - 15C_1$.

$$\begin{bmatrix} 3 & 4 & 0 \\ 7 & 9 & 0 \\ 13 & 2 & 0 \end{bmatrix}$$

$\therefore$ Determinant of the matrix $= 0$.

(3) If the matrix A is such that

$$A = \begin{bmatrix} 2 \\ -4 \\ 7 \end{bmatrix} \begin{bmatrix} 1 & 9 & 5 \end{bmatrix}$$

$$A = \begin{bmatrix} 2 & 18 & 10 \\ -4 & -36 & -20 \\ 7 & 63 & 35 \end{bmatrix}$$

$$C_3 = 5 \times C_1$$

$\therefore |A| = 0$.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

(4) $\begin{vmatrix} 1 & x & x^2 \\ 1 & y & y^2 \\ 1 & z & z^2 \end{vmatrix}$.

Which of the following is the matrix determinant NOT equal to

A $\begin{vmatrix} 1 & x(x+1) & (x+1) \\ 1 & y(y+1) & (y+1) \\ 1 & z(z+1) & (z+1) \end{vmatrix}$

B $\begin{vmatrix} 1 & x+1 & x^2+1 \\ 1 & y+1 & y^2+1 \\ 1 & z+1 & z^2+1 \end{vmatrix}$

C. $\begin{vmatrix} 0 & x-y & x^2-y^2 \\ 0 & y-z & y^2-z^2 \\ 1 & z & z^2 \end{vmatrix}$.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

D. $\begin{vmatrix} 2 & x+y & x^2+y^2 \\ 2 & y+z & y^2+z^2 \\ 1 & z & z^2 \end{vmatrix}$

Soln.

Clearly by seeing option B if we perform $C_2 \to C_2 - C_1$

$C_3 \to C_3 - C_1$

we get back the orignl Matrix.

Similarly for Matrix of option C

if we perform $R_2 \leftarrow R_2 + R_3$

$R_1 \leftarrow R_1 + R_2$.

We get back the original matrix.

Also on option D. if we observe $C_2 \leftarrow C_2 - C_3$

$C_3 \leftarrow C_3 - C_1$

We get back the original matrix, the only remaing option is option 1.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

(5) Let A, B, C, D be $n \times n$ matrices, each with non-zero determinant. If $ABCD = I$, then B^{-1} is .

A. $B^{-1} C^{-1} A^{-1}$

B. CDA

C. ADC.

D. Does not necessarily exist.

Soln

$$ABCD = I.$$

$$ABCD * D^{-1} = I \cdot D^{-1}$$

$$ABC * C^{-1} = D^{-1} C^{-1}$$

$$A^{-1} \cdot AB = A^{-1} B^{-1} C^{-1}$$

$$\Rightarrow B = A^{-1} D^{-1} C^{-1}$$

$$(AB)^{-1} = B^{-1} A^{-1}$$

$$\therefore B^{-1} = CDA$$

option is B.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

PH: +91-844-844-0102

1.12 Special Types of Matrices and Properties

Orthogonal Matrix :-

- It is a square matrix
- $Q \cdot Q^T = I$ or $Q^T = Q^{-1}$

Intuitive Understanding

$$\begin{bmatrix} \leftarrow r_1 \rightarrow \\ \leftarrow r_2 \rightarrow \\ \leftarrow r_3 \rightarrow \end{bmatrix} \begin{bmatrix} \uparrow & \uparrow & \uparrow \\ r_1 & r_2 & r_3 \\ \downarrow & \downarrow & \downarrow \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Q $\qquad$ Q^T

$r_1^T \cdot r_1 = 1$

$r_1 \cdot r_2 = r_1^T r_2 = 0$

$r_2^T r_2 = 1$

$r_i \cdot r_j = 0$ if $i \neq j$

$r_i \cdot r_j = 1$ if $i = j$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

r_i's are unit vector.

every r_i and r_j if $i \neq j$ are perpendicular vectors

and $i = j$ are parallel vectors.

- If we think of it from the perspective of a linear transform

$Q =$ 

3×3.

$\hat{i}, \hat{j}, \hat{k}$

Initially $\hat{i} \perp \hat{j} \perp \hat{k}$,

on transformation

$\hat{i}, \perp \hat{j}, \perp \hat{k},$

$\| \hat{i}, \| = \| \hat{j}, \| = \| \hat{k} \| = 1$

→ Rotation with no shear and no scaling is one example of an orthogonal matrix.



→ Mirror or Reflection is an example of an orthogonal matrix.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

→ $|Q| = \pm 1$ volume of the parallelopiped $= \pm 1$.

volume does not change for an area only. orientation will change.

Preservation of Dot Product.

→ $U.V = U^T V = (Q.U).(QV)$

$= (QU)^T (QV)$

$= U^T Q^T Q V$

$\Rightarrow \underline{U^T V}$.

- If the rows of an orthogonal matrix are interchanged it still remains an orthogonal matrix.

IDEMPOTENT MATRIX.

- It is a square Matrix such that $M \times M = M$

$$M = \begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{bmatrix}$$

$M = I$

$|M| = 0$ or

Gatecse@appliedcourse.com

Gate.Appliedcourse.com

Ph: +91 844-844-0102

- A is idempotent $A^m = A$ for $m = 1, 2, \ldots$

Applications in ML & AI.

Involutary Matrix

$A^2 = I$ (Square Root Matrix of I).

or $A = A^{-1}$.

Geometrically :- If A is a linear tranformation and its inverse is equal to itself then A is an involutary Matrix

Ex

$$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \qquad i = \sqrt{-1}$$

$A^2 = I$ & A is symmetric $\longrightarrow$ A is orthogonal.

$A \cdot A = I$ & $A = A^T$ $\longrightarrow$ $A \cdot A^T = I$.

- An another way to look at it geometrically is an involutary Matrix is like a mirror reflection, if applied twice we get back the orginal image/figure.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

Properties of An Involutary Matrix

① $|A| = \pm 1$

② $\frac{1}{2}(A+I)$ is idempotent.

③ If A and B are Involutary then. AB is also involutary.

④ If A is involutary $A^n = \begin{cases} A & \text{if } n \text{ is odd} \\ I & \text{if } n \text{ is even.} \end{cases}$

NILPOTENT MATRIX

If $N^K = 0$ for some K = +ve integer

Smallest K is index of N.

ex $\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ $A^2 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$

Any triangular matrix with diagonal elements = 0 is a nilpotent matrix. with (index $< n$)

$\begin{bmatrix} 0 & 1 & 2 \\ 0 & 0 & 3 \\ 0 & 0 & 0 \end{bmatrix}$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

Properties of Nilpotent Matrix.

(1) $|I + N| = 1$

(2) $(I+N)^{-1} = \underbrace{1 - N + N^2 - N^3 + \dots}_{} + \underbrace{N^R + N^{R+1} + \dots}_{0}$

as $N^R = 0$

$N^{R+1} = 0$.

. .

1.13 Trace Of A Matrix

Equality Of Matrices

$A_{m \times m} = B_{m \times n}$

- Orders should be same

and $A_{ij} = B_{ij} \quad \forall i,j$.

Trace Of a Matrix

$$\text{trace}(A) = \sum_{i=1}^{n} a_{ii}$$

$$\text{trace}\left(\begin{bmatrix} -1 & 0 & 3 \\ 11 & 5 & 2 \\ 6 & 12 & -5 \end{bmatrix}\right) = -1 + 5 - 5 = -1$$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

Geometric Intuition :- Related to Eigen values and Eigen Vectors to be covered later.

→ $Trace(A+B) = Trace(A) + Trace(B)$.

→ $Trace(c.A) = c.Trace(A)$.

→ $Trace(A.B) = Trace(B.A)$ As long as AB and BA are defined.

→ $Trace(AB^T) = Trace(A^TB) = Trace(BA^T) = Trace(B^TA)$.

→ Cyclic Property of Trace :

$Trace(ABCD) = Trace(BCDA) = Trace(CDAB) = Trace(DABC)$

→ If A, B, C are symmetric

$Trace(ABC) = Trace(CBA) = Trace(ACB) = Trace(CAB)$.

$= Trace(BAC) = Trace(BCA)$.

→ Common Mistake

$trace(AB) \neq Trace(A).Trace(B)$.

→ $trace(P^{-1}AP) = trace(P^{-1}(AP))$

$= trace(P^{-1}PA)$

$= trace(A)$.

Gatecse@appliedcourse.com

GateAppliedcourse.com

Ph: +91-844-844-0102

- If A is a symmetric Matrix; B is a skew symmetric matrix.

trace (AB) = 0 -

1.14 Rank Of A Matrix

→ Very useful for solving a system of linear equations.

→ Rank is defined for both a square and rectangular matrix.

$$A = \begin{bmatrix} 1 & 2 \\ -1 & -2 \end{bmatrix} \qquad C_1 = \frac{C_2}{2}$$

$\hat{i}$, lands $\hat{j}$, lands.

If we perform transformation for any vector $\vec{v}$.

$$\underset{2\times 2}{A}\ \underset{2\times 1}{V} = \underset{2\times 1}{U}$$



Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph.: +91 814-814-0102

- The transformed vector will lie on the same line as $\hat{i}$, and $\hat{j}$, lie on the same line

$|A| = 0$ area of the parallelogram = 0. as $\hat{i}$, and $\hat{j}$, are linearly dependent.

Geometric intuition of Rank(A).

- The dimension of the vector space spanned by the column vectors of A span.

$$A = \begin{matrix} C_1 & C_2 & C_3 \end{matrix} \begin{bmatrix} 1 & 0 & 1 \\ -2 & -3 & 1 \\ 3 & 3 & 0 \end{bmatrix}$$

Alternate way of thinking

Rank(A) = Maximum no of linearly independent columns of A.

If $C_1 \longleftarrow a C_2 + b C_3$ C_1 is linearly dependent on C_2 and C_3.

Numerical way to think about Rank(A)

If A is a square matrix

if $|A| = 0$ Rank $\neq n$, Rank $< n$ } — Rank(A) = size of largest sub matrix with non zero determinant.

$|A| \neq 0$ Rank(A) = n

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

→ Column Rank is what we have discussed so far. based on column

⇒ Row Rank of a matrix is defined based on the rows of the matrix

- Max no of linearly independent rows of that matrix = Rank.

- Column Rank of the Matrix = Row rank of the matrix = Rank of the Matrix.

→ $Rank(A) = Rank(A^T)$

—

example

$$A = \begin{bmatrix} 1 & 1 & 0 & 2 \\ -1 & -1 & 0 & -2 \end{bmatrix} \begin{matrix} \leftarrow R_1 \\ \leftarrow R_2 \end{matrix}$$

$R_1 = -1 * R_2$.

Rank = 1.

→ If we have an mxn matrix A.

$Rank(A) \leq \min(m, n)$.

Compute Rank

→ Elementary Row operations do not change the rank of a matrix.

1. Swap R_i and R_j

2. $R_i \leftarrow cR_i$ $c \neq 0$

3. $R_i \leftarrow R_i + kR_j$.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Gauss Elimination Method

Ph: +91 844-844-0102

$$\begin{bmatrix} 1 & 3 & 1 & 9 \\ 1 & 1 & -1 & 1 \\ 3 & 11 & 5 & 35 \end{bmatrix} \xrightarrow[R_3 \leftarrow R_3 - 3R_1]{R_2 \rightarrow R_2 - R_1} \begin{bmatrix} 1 & 3 & 1 & 9 \\ 0 & -2 & -2 & -8 \\ 0 & 2 & 2 & 8 \end{bmatrix}$$

$$R_3 \rightarrow R_3 + R_2 \cdot \begin{bmatrix} 1 & 3 & 1 & 9 \\ 0 & -2 & -2 & -8 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

# of non zero rows = Rank of the Matrix. = 2

The above form is the Echelon form of the Matrix.

Echelon Form

- The no of 0's before the first non zero number in any row is less than the no of such zeros in the next row.
- All zero rows must be below non zero rows.

(1) $\text{rank}(A) \leq \min(m, n)$

(2) Only zero matrix have rank = 0.

(3) full row rank = rank = m.
full column rank = rank = n.

Gatecse@appliedcourse.com

Gate.Appliedcourse.com

Ph: +91 844-844-0102

$A: n \times n$ full rank $\rightarrow$ rank $= n$

(4) $A_{n\times n}$ is invertable iff Rank $(A) = n$.

if rank $(A) < n$ A is not invertable.

(5) $\text{Rank}(AB) \leq \text{Min}(\text{Rank}(A), \text{Rank}(B))$.

(6) $\text{Rank}(A+B) \leq \text{Rank}(A) + \text{Rank}(B)$

(7) Sylvester's Rank inequality.

$$\text{rank}(A) + \text{rank}(B) - n \leq \text{rank}(AB).$$

(8) $\text{rank}(A^TA) = \text{rank}(AA^T) = \text{rank}(A) = \text{rank}(A^T)$

(9)

$$P = \begin{bmatrix} 1 & 1 & -1 \\ 2 & -3 & 4 \\ 3 & -2 & 3 \end{bmatrix} \qquad Q = \begin{bmatrix} -1 & -2 & -1 \\ 6 & 12 & 6 \\ 5 & 10 & 5 \end{bmatrix}$$

$$\text{Rank}(P+Q) = \begin{bmatrix} 0 & -1 & -2 \\ 8 & 9 & 10 \\ 8 & 8 & 8 \end{bmatrix} \xrightarrow{R_2 \leftarrow R_2 - R_3}$$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

$$\begin{bmatrix} 8 & 8 & 8 \\ 0 & 1 & 2 \\ 0 & -1 & -2 \end{bmatrix} \xrightarrow{R_3 \leftarrow R_3 + R_2} \begin{bmatrix} 8 & 8 & 8 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$$

$\text{Rank}(A) = 2$

en2

$$\begin{bmatrix} 1 & 4 & 8 & 7 \\ 0 & 0 & 3 & 0 \\ 4 & 2 & 3 & 1 \\ 3 & 12 & 24 & 21 \end{bmatrix} \xrightarrow{R_2 \leftarrow R_3} \begin{bmatrix} 1 & 4 & 8 & 7 \\ 4 & 2 & 3 & 1 \\ 0 & 0 & 3 & 0 \\ 3 & 12 & 24 & 21 \end{bmatrix}$$

$R_2 \leftarrow R_2 - 4R_1 \qquad R_3 \leftarrow R_3 - 3R_1$

$$\begin{bmatrix} 1 & 4 & 8 & 7 \\ 0 & -14 & -29 & -27 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Rank = 3

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph. +91 844-844-0102

1.15 Systems of linear Equations - 1

- V. important application of linear Algebra.

$$3x + 2y - z = 1$$
$$2x - 2y + 4z = -2$$
$$-x + \frac{y}{2} - z = 0.$$

↓

$$\begin{bmatrix} 3 & 2 & -1 \\ 2 & -1 & 4 \\ -1 & \frac{1}{2} & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ -2 \\ 0 \end{bmatrix}$$

Coeff Matrix A x b.

$$Ax = b.$$

Geometric interpretation

1. $Ax = b$.
   ↓ i/p ↓ o/p.
   ↓ transformation

For the linear transformation represented by A and output b what is the input x.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 8448440102

② Wikipedia https://en.wikipedia.org/wiki/System_of_linear_equations

→ Canonical form of eqn in 2D = $ax+by+c=0$ - st line.

→ In 3D coordinate geometry → $ax+by+cz+d=0$ represents a plane in 3D.

→ In 4D space the equation $ax+by+cz+dt+e=0$ - represents a hyperplane in 4D.

→ Similarly a linear equation in 10 variables represents a hyperplane in 10D space.

→ 3 Equations represents 3 planes in 3D space and the solution represented is the pt of intersection of the 3 planes.

Solution Set :- Set of all valid solutions to the linear system of equations

1. Infinitely many solutions → l_1, l_2 are lapping lines
2. Single unique solution → l_1 l_2
3. No Solutions. → l_1 // l_2 parallel lines.

They will fall in one of the three categories.

Underdetermined and Over determined System of Equations

n - no of variables m - no of equations

① $m<n$ - Underdetermined System (Cannot determine the soln)

② $m=n$ - Exactly determined System. (1 soln, no soln, ∞ solutions)

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

③ Over determined system of equations. ($m > n$).

We could have more than one solutions



$n=2$ $m=3$

Independence of System of Linear Equations :- A linear system of eqns is said to be independent if none of the equations can be arrived at from other equations algebraically.

ex1

① $3x + 6y = 6$.

② $6x + 4y = 12$

② = 2 * ①.

linearly dependent system of Equations.

ex2

① $x - 2y = -1$

② $3x + 6y = 8$.

③ $4x + 3y = 7$.

③ = ① + ②

linearly dependent on system of Equation.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

Inconsistent System of Equations

- If the system of equations is inconsistant :- No solution exists



|| lines.

- If the coefficient Matrix rows are linearly dependent and the constants do not satisfy the linear dependence then the given system of equations is inconsistent.

$$3x+2y=12$$
$$3x+2y=6.$$

$$\begin{bmatrix} 3 & 2 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 12 \\ 6 \end{bmatrix}$$

$R_1 = R_2 \times 1$

But in the constant matrix the same linear dependence does not hold.

Ex-2

$$\begin{bmatrix} 3 & 2 \\ 6 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 12 \\ 6 \end{bmatrix}$$

$R_1 = \frac{1}{2} R_2$ and also in the constant matrix the same holds

It means that the two lines are overlapping and the system of

Gate.Appliedcourse.com
Gatecse@appliedcourse.com

equations has infinite no of solutions.

→ If the rows of the coff. Matrix are linearly then the system of linear Equations is consistant.

Summary.



System Of linear Equations -2.

Solving System of L. Equations

Elimination of Variables.

$x + 3y - 2z = 5$ -1 ⟶ $x = 5 + 2z - 3y$.

$3x + 5y + 6z = 7$ -2

$2x + 4y + 3z = 8$ -3

substition in eq 2 and 3

Ph. +91 844-844-0102
Gate.Appliedcourse.com
Gatecse@appliedcourse.com

Ph: +91 844-844-0102

$-4y + 12z = -8 \Rightarrow y = 2 + 3z$ — (4)

Substitute in eq 3.

we get $z = 2$

Substitute in eq (4)

We get $y = 8$

Substitute the values of y and z in any eq 1, 2, or 3 we get $x = -15$

GAUSS ELIMINATION METHOD.

The coefficients of the equation and the constants are written in a single matrix known as augmented matrix

$$\left[\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 3 & 5 & 6 & 7 \\ 2 & 4 & 3 & 8 \end{array}\right]$$

A | b.

Need to reduce to 0.

$R_2 \longrightarrow R_2 - 3R_1$

$R_3 \longrightarrow R_3 - 2R_1$

$$\sim \left[\begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & -4 & +12 & -8 \\ 0 & -2 & 7 & -2 \end{array}\right]$$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

$R_3 \longrightarrow R_3 - \frac{1}{2} R_2$.

$$\begin{bmatrix} 1 & 3 & -2 & 5 \\ 0 & -4 & 12 & -8 \\ 0 & 0 & -2 & -4 \end{bmatrix}$$

$$-2z = -4$$

$$z = 2$$

We can substitute in the 2nd row.

$$-4y + 12(z) = -8.$$

$$-4y + 24 = -8$$

$$y = 8.$$

Now we can substitute values of y and z in R_1

$$x + 3y - 2z = 5$$

$$x + 3(8) - 2(2) = 5.$$

$$x + 24 - 4 = 5$$

$$x = -15$$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph:+91 844 844 0102

→ Another way to solve is to proceed further and reduce the matrix in the form of an identity matrix, then the constants represent the values of x, y and z.

③ Cramers Rule.

$$1x + 3y - 2z = 5$$
$$3x + 5y + 6z = 7$$
$$2x + 4y + 3z = 8.$$

$$Ax = b.$$

constants

$$x = \frac{\begin{vmatrix} 5 & 3 & -2 \\ 7 & 5 & 6 \\ 8 & 4 & 3 \end{vmatrix}}{\begin{vmatrix} 1 & 3 & -2 \\ 3 & 5 & 6 \\ 2 & 4 & 3 \end{vmatrix}} \qquad y = \frac{\begin{vmatrix} 1 & 5 & -2 \\ 3 & 7 & 6 \\ 2 & 8 & 3 \end{vmatrix}}{\begin{vmatrix} 1 & 3 & -2 \\ 3 & 5 & 6 \\ 2 & 4 & 3 \end{vmatrix}}$$

constants.

→ Not used much in practice as the calculation of determinants is not computationally efficient

However a geometric intuition of Cramers rule is available on a youtube video of 3 Blue 1 brown channel.

https://youtu.be/jBSC34PxzoM.

Gatecse@appliedcourse.com

Gate.Appliedcourse.com

PH: +91 844-844-0102

④ Matrix Solution way to Solve System of linear Equations :-

$$Ax = b.$$

$$x = A^{-1}b.$$

→ If $|A|=0$ - A is not invertable ⇒ ∞ no of solutions
or
No solutions.

→ Easiest way amongall or most preffred is Cramers Elimination method.

Homogenous system of linear Equations

$$Ax = b$$

if $b = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \rightarrow Ax = 0$

$3x + 2y.$

$x + 2y = 0.$

$x = 0 \quad y = 0.$

① Zero/trivial soln $= \begin{bmatrix} 0 \\ 0 \end{bmatrix}$.

② If $|A| \neq 0$ then only zero solution exists.

③ If $|A| = 0$ then ∞- many solution exists.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 8448440102

(4) u & v are solution of $Ax=0$ then $u+v$ is also a solution.

$$A(u+v) = Au + Av = 0$$

(5) u is a solution $\longrightarrow$ cu is also a solution

(6) $Ax = b$ $\qquad$ $Ax = 0$.

$\uparrow$ $\qquad$ $\uparrow$

p is a soln $\qquad$ v is a solution

Then $p+v$ is also a soln to ~~Ax=b~~ $Ax = b$.

1.17 Solved Problems GATE 2017, 2014, 2015, 2005, 2004

Q Let $c_1, c_2, \ldots c_n$ be scalars not all zero, such that the following expression holds.

$$\sum_{i=1}^{n} c_i a_i = 0 \cdot \quad \text{where } a_i \text{ is the column vector of } \mathbb{R}^n$$

Consider the set of linear Equations

$$Ax = B$$

where $A = [a_1 a_2 \ldots a_n]$ and $b = \sum_{i=1}^{n} a_i$

Then the sol of Equations has.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: 91-844-844-0102

A. A unique solution, has $x = j_n$ where j denotes n dimensional vector for all i.

B. No solution.

C. Infinitely many solutions

D. Finitely many solutions

Solution.

given $\sum_{i=1}^{n} c_i a_i = 0$.

c_i's are constants and a_i's represent the columns of the matrix

which mean in case of a 3×3 matrix

$$c_1 a_1 + c_2 a_2 + c_3 a_3 = 0.$$

$$a_1 = -\frac{c_2}{c_1} a_2 - \frac{c_3}{c_1} a_3$$

a_1 is linearly dependent on a_2 and a_3.

Now this means that $|A| = 0$, the system of equations either have infinitely many solutions or no solutions.

Also we are given that $\sum_{i=1}^{n} a_i = b$.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

Which means that the sum of coffients = constant matrix.

i.e in place of x, y, z we put (1, 1, 1) we will get the sums of the coffients, in other words (1, 1, 1) is a solution to the system of linear equations, we cannot say that no Soln exist, and therefore the system has infinite no of solns. option C.

⟶ 8 Consider a system, each consisting of m linear equations in n variables.

I. If $m < n$ then all such systems have a solution.

II. If $m > n$, then none of these systems has a solution.

III. If $m = n$ then there exist a system which has a solution

Which of the following is correct?

(A) I, II and III are true

(B) Only II and III are true

(C) Only III is true

(D) None of them is true.

Soln

Statement I. Lets take $m = 1$ $n = 2$.

This is an underdetermined system, all such systems do not have a soln it just represents one line.

Gatecse@appliedcourse.com

Gate.Appliedcourse.com

Ph: +91 844-844-0102

Statement II

$m=3, n=2$.

↓.

3 lines

L_2, L_4





Soln may exist, Statement II is false.

Statement III. If $m=n$

$m=2\ n=2$.

2 lines



soln exists



does

Solutions ~~need~~ not exist.

∴ Statement 3 is true.

option c is correct.

8

→ If the following system has a non trivial solution

$$px + qy + rz = 0$$

$$qx + ry + pz = 0$$

$$rx + py + qz = 0.$$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Which of the following options is True?

(A) $p - q + r = 0$ or $p = q - r$.

(B) $p + q - r = 0$ or $p = -q = r$.

(C) $p + q + r = 0$ or $p = q = r$.

(D) $p - q + r = 0$ or $p = -q = -r$.

Sol: If a homogeneous sys of linear equations then $|A| = 0$

$$\begin{vmatrix} p & q & r \\ q & r & p \\ r & p & q \end{vmatrix} = 0.$$

If we ~~opt~~ observe the 3rd option $p = q = r$ $|A| = 0$ option 3 is correct

Q → Consider the following system of equations in three real numbers.

x_1, x_2, x_3.

$$2x_1 - x_2 + 3x_3 = 1$$

$$3x_1 - 2x_2 + 5x_3 = 2$$

$$-x_1 + 4x_2 + x_3 = 3.$$

Ph: +91 844-844-0102
Gate.Appliedcourse.com
Gatecse@appliedcourse.com

Ph: +91 844-844-0102

The system of equations have

(A) No solutions

(B) A Unique Solution

(C) More than one but finite no of solutions

(D) An infinite no of solutions.

$$A = \begin{bmatrix} 2 & -1 & 3 \\ 3 & -2 & 5 \\ -1 & 4 & 1 \end{bmatrix}$$

$|A| = -6 \neq 0$ ∴ The system has a unique soln option B.

Q → What values of x, y, z would satisfy the following system of linear Equations

$$\begin{bmatrix} 1 & 2 & 3 \\ 1 & 3 & 4 \\ 2 & 2 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 6 \\ 8 \\ 12 \end{bmatrix}$$

A) $x=6, y=3, z=2$

B) $x=12\ y=3\ z=-4$

C) $x=6\ y=6\ z=-4$

D) $x=12\ y=-3\ z=0$.

Soln on Substituting values we can come to a solution

option C is correct.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

PH: +91 844-844-0102

1.18 Eigen Values And Eigen Vectors

Geometric intution

$A_{n \times n}$ → Square Matrix {Transformation in nD space}

$$Av = \lambda v$$

$v \in \mathbb{R}^d$. real valued d-dim vector.

v is the eigen vector of A

λ is the eigen value of A.

→ v is that special vector of A if we apply linear tranformation of A it is equalent to scaling it by a factor of λ.



$|\lambda| > 1$ ⇒ stretched

$|\lambda| < 1$ ⇒ shrunk.

$|\lambda| < 0$ reversed.

Gate Applied course.com

Gatecse@appliedcourse.com

Animations from 3Blue1Brown youtube channel for Eigen value and Eigen Vectors :- http://youtu.be/PFDu9oVAE-g.

Ph: +91 844-844-0102

Algebraic and Numeric way

$$AV = \lambda V$$

$$\Rightarrow AV = \lambda IV$$

$$\Rightarrow AV - \lambda IV = 0.$$

$$\Rightarrow (A - \lambda I)V = 0.$$

If $V \neq 0$.

$|A - \lambda I| = 0 \rightarrow$ Characteristic Eqn.

$$A = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$$

$$|A - \lambda I| = 0 \Rightarrow \left| \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} \right|$$

$$\Rightarrow \begin{vmatrix} 2-\lambda & 1 \\ 1 & 2-\lambda \end{vmatrix} = (2-\lambda)^2 - 1 = 0.$$

$$\lambda^2 - 4\lambda - 3 = 0.$$

$$\lambda_1 = 3, \lambda_2 = 1$$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

$(A-\lambda I)V_1 = 0$.

$\lambda_1 = 3$.

$$\begin{bmatrix} 2 & -3 \\ 1 & 2-3 \end{bmatrix} \begin{bmatrix} V_{11} \\ V_{12} \end{bmatrix} = 0.$$

$$-V_{11} + V_{12} = 0$$

$$V_{11} - V_{12} = 0.$$

$$V_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$\lambda_2 = 1 \longrightarrow V_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$



Properties of E.V. & E.Vectors

① It is possible that A has no eigen value and eigen vectors

ex Rotation Matrix $A_R = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = R_{90^\circ}$.

∴ Such a matrix does not have any eigen values

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

(2) $A = S = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$



$|A - \lambda I| = 0$

$(2-\lambda)(2-\lambda) = 0$.

$\lambda = 2$ single value.

No of eigen values $<= n$ for a $n \times n$ matrix.

(3) A = Diagonal Matrix $= \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$ $\lambda = 2, 3$.

$(2-\lambda)(3-\lambda) = 0$.

→ Diagonal Matrix are the E. values. and basis vectors are the eigen vectors.

(4) $\text{Trace}(A) = \sum_{i=1}^{n} \lambda_i$

(5) $|A| = \prod_{i=1}^{n} \lambda_i$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-8440102

(8) If λ is the eigen value of A then λ^n is is eigen value of A^n.

2. $K\lambda$ is eigen value of (KA)

K is a scalar $\neq 0$

3. $\lambda + K$ = Eigen value of $(A+KI)$

9 → Eigen value of the matrix represented by $a_0 I + a_1 A + a_2 A^2$ is $a_0 + a_1 \lambda + a_2 \lambda^2$.

(9) If λ is eigen value of A and $|A| \neq 0$ then $\frac{1}{\lambda}$ is eigen value of A^{-1}.

(10) Eigen value of $(Adj(A)) = \frac{|A|}{\lambda}$.

(11) In case of a triangular matrix the eigen values are the digonal elements itself.

$$\begin{bmatrix} 1 & 0 & 0 \\ 1 & 2 & 0 \\ 2 & 3 & 3 \end{bmatrix} \qquad (1-\lambda)(2-\lambda)(3-\lambda) = 0$$

$$\lambda = 1, 2, 3$$

$$\begin{bmatrix} 2 & 3 & 0 \\ 0 & 6 & 1 \\ 0 & 0 & 7 \end{bmatrix} \rightarrow \lambda = 2, 6, 7.$$

True in case of upper and lower tregular matrix.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

(12) Eigen Value(A) = Eigen Value(A^T)

→ If A is an orthogonal matrix then $AA^T = I$ $A^T = A^{-1}$

λ is an eigen value of A then $\frac{1}{\lambda}$ is also an eigen value of A.

(13) If we have a real and symmetric matrix, eigen values are real valued.

→ if $a + ib$ is an eigen value of A then $a - ib$ is also an eigen value of A.

(14) If For a skew symmetric matrix A which is real valued, the eigen values are Zero or Purely Imaginary.

(15) Eigen Value (AB) = Eigen Value (BA).

(16) If $\lambda_1, \lambda_2 \ldots \lambda_n$ are non repeating eigen values of A then the eigen vectors of A $V_1, V_2 \ldots V_n$ are linearly independent.

Gate Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844 844-0102

1.9 Cayley-Hamilton Theorem And Diagnolization

- If we have a matrix A, its characteristic polynomial is given by.

$$P(\lambda) = |A - \lambda I| = 0.$$

For example $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ $P(\lambda) = |A - \lambda I| = \lambda^2 - 5\lambda - 2 = 0.$

→ The matrix itself satisfies the characteristic Equation. (Cayley Hamilton theorem)

$$A^2 - 5A - 2I = 0.$$

Applications

1. We can represent higher powers of A in terms of simple operations on matrix A. we can do this very fast this way.

$$A^2 = 5A + 2I$$

$$A^3 = 5A^2 + 2A$$

$$= 5(5A + 2I) + 2A$$

$$= 27A + 10I$$

Similarly we can compute $A^4, A^5 \ldots$ any power of A.

2. Compute inverse.

$$A^2 = 5A + 2I$$

Multiply by A^{-1} throughout. $\Rightarrow$ $A = 5I + 2A^{-1}$

$$\Rightarrow A^{-1} = \frac{A}{2} - \frac{5}{2}I$$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Also higher - ve exponents can be calculated by Cayley Hamilton theorem.

Ph: +91 844-844-0102

Diagonalization : A matrix "A" is said to be diagonalizable if there exists a matrix P such that $P^{-1}AP$ = diagonal matrix.

$$\begin{bmatrix} a_1 & \cdots & 0 \\ 0 & a_2 & \cdots \\ \vdots & \vdots & \ddots \\ 0 & \cdots & a_n \end{bmatrix}$$

If we write the equation

$$P^{-1}AP = \begin{bmatrix} \lambda_1 & 0 & 0 & \cdots \\ 0 & \lambda_2 & 0 & \cdots \\ 0 & 0 & \lambda_3 & \cdots \\ \vdots & \vdots & \vdots & \ddots \\ \cdots & \cdots & 0 & \lambda_n \end{bmatrix}$$

Multiplying both sides with P.

$$AP = P \begin{bmatrix} \lambda_1 & 0 & 0 & \cdots \\ 0 & \lambda_2 & 0 & \\ 0 & 0 & \lambda_3 & \cdots \\ 0 & \cdots & & \\ \cdots & \cdots & \cdots & \lambda_n \end{bmatrix}$$

If $P = \begin{bmatrix} \uparrow & \uparrow & \uparrow & & \uparrow \\ V_1 & V_2 & V_3 & \cdots & V_n \\ \downarrow & \downarrow & \downarrow & & \downarrow \end{bmatrix}$ $n \times n$.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

$$A \cdot P = \begin{bmatrix} \uparrow & \uparrow & & \uparrow \\ AV_1 & AV_2 & \cdots & AV_n \\ \downarrow & \downarrow & & \downarrow \end{bmatrix} = \begin{bmatrix} \uparrow & \uparrow & \uparrow & \uparrow \\ V_1\lambda_1 & V_2\lambda_2 & V_3\lambda_3 & \cdots V_n\lambda_n \\ \downarrow & \downarrow & \downarrow & \downarrow \end{bmatrix}$$

$$\therefore \quad \begin{aligned} AV_1 &= V_1\lambda_1 \\ AV_2 &= V_2\lambda_2 \\ &\vdots \\ AV_n &= V_n\lambda_n \end{aligned}$$

} This is the definition of Eigen Values, Vectors.

Given a matrix A we can easily find the diagonalization matrix by using its eigen vectors.

$$P = \begin{bmatrix} \uparrow & \uparrow & & \uparrow \\ V_1 & V_2 & \cdots & V_n \\ \downarrow & \downarrow & & \downarrow \end{bmatrix}$$

$$P^{-1}AP = \begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & \cdots \\ \cdots & \cdots & \lambda_n \end{bmatrix} = D$$

→ *. For the matrix A to be diagonalizable the matrix P should be invertible i.e in other words the Eigen Vectors should be Linearly Independent.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

$A = \begin{bmatrix} 1 & 2 & 0 \\ 0 & 3 & 0 \\ 2 & -4 & 2 \end{bmatrix}$ on calculating E.V we get $\lambda_1 = 3$ $\lambda_2 = 2$ $\lambda_3 = 1$.

$V_1 = \begin{bmatrix} -1 \\ -1 \\ 2 \end{bmatrix}$ $V_2 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ $V_3 = \begin{bmatrix} -1 \\ 0 \\ 2 \end{bmatrix}$

$$P = \begin{bmatrix} -1 & 0 & -1 \\ -1 & 0 & 0 \\ 2 & 1 & 2 \end{bmatrix}$$

$$P^{-1} = \begin{bmatrix} 0 & -1 & 0 \\ 2 & 0 & 1 \\ -1 & 1 & 0 \end{bmatrix}$$

We get.

$$P^{-1}AP = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Applications of diagonalization

→ Compute Higher Exponents of any matrix.

$P^{-1}AP = D.$

$PP^{-1}APP^{-1} = PDP^{-1}$

$A^2 = PDP^{-1}PDP^{-1}$

$\Rightarrow PD^2P^{-1}$

Gate-Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

Similarly $A^k = PD^kP^{-1}$.

$$A^{100} = PD^{100}P^{-1}$$

Instead of 100 matrix multiplications we can do just 2 matrix multiplications

As it is very easy to compute exponent of diagonal Matrices.

1.20 LU Decomposition

- Very similar to Gauss Elimination
- If we have a system of linear equations $Ax=b$ and we have to find a solution to it i.e. we need to find x.
- We try to decompose A as $A=LU$ where L is a lower triangular matrix and U is an upper triangular matrix.

For example lets consider $A = \begin{bmatrix} 1 & 2 & 4 \\ 3 & 8 & 14 \\ 2 & 6 & 13 \end{bmatrix}$ if we are trying to decompose A as LU

$$L = \begin{bmatrix} 1 & 0 & 0 \\ L_{21} & 1 & 0 \\ L_{31} & L_{32} & L_{33} \end{bmatrix} \quad U = \begin{bmatrix} U_{11} & U_{12} & U_{13} \\ 0 & U_{22} & U_{23} \\ 0 & 0 & U_{33} \end{bmatrix}$$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

On multiplying L and U we get

$$LU = \begin{bmatrix} U_{11} & U_{12} & U_{13} \\ L_{21}U_{11} & (L_{21}U_{12}+U_{22}) & (L_{21}U_{13}+U_{23}) \\ L_{31}U_{11} & (L_{31}U_{12}+L_{32}U_{22}) & \begin{pmatrix} L_{31}U_{13}+ \\ L_{32}U_{23}+ \\ U_{33} \end{pmatrix} \end{bmatrix}$$

On comparing the above matrix with A and equating the corresponding terms

$U_{11} = 1$

$U_{12} = 2$

$U_{13} = 4$

$L_{21}U_{11} = 3$

$L_{21}(1) = 3 \Rightarrow L_{21} = 3$

$L_{21}U_{12} + U_{22} = 8$

$3(2) + U_{22} = 8$

$U_{22} = 2$

$L_{21}U_{13} + U_{23} = 14$

$3(4) + U_{23} = 14$

$U_{23} = 2$

$L_{31}U_{11} = 2$

$L_{31} = 2$

$L_{31}U_{12} + L_{32}U_{22} = 6$

$2(2) + L_{32}(2) = 6$

$L_{32} = 1$

$L_{31}U_{13} + L_{32}U_{23} + U_{33} = 13$

$2(4) + 1(2) + U_{33} = 13$

$U_{33} = 3$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

$$A = \begin{bmatrix} 1 & 2 & 4 \\ 3 & 8 & 14 \\ 2 & 6 & 13 \end{bmatrix} = LU = \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 2 & 1 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 2 & 4 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{bmatrix}$$

We can use LU Decomposition to solve system of linear Equations.

Step 1 :- Decompose the coeff Matrix A into two lower and upper triangular matrix.

Step 2 :-

$$LUx = b$$

$$\downarrow$$

$$y$$

$$Ly = b$$

solve for y.

$$\begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 2 & 1 & 1 \end{bmatrix} y = \begin{bmatrix} 3 \\ 13 \\ 4 \end{bmatrix} \qquad y = \begin{bmatrix} 3 \\ 4 \\ -6 \end{bmatrix}$$

Step 3 Now solve for x.

$$Ux = y$$

$$\begin{bmatrix} 1 & 2 & 4 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 3 \\ 4 \\ -6 \end{bmatrix}$$

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

→ This method is very easy to solve using the LU decomposition as we have upper and lower triangular matrices and they are already in reduced form.

Ph: +91 844-844-0102

→ Does Every matrix have a LU decomposition? - No

We need to check the determinant of the leading matrices.

$$A = \begin{bmatrix} 1 & 2 & 4 \\ 3 & 8 & 14 \\ 2 & 6 & 13 \end{bmatrix}$$

$A_1 = 1$

$|A_1| = 1 \neq 0$

$$A_2 = \begin{bmatrix} 1 & 2 \\ 3 & 8 \end{bmatrix}$$

$|A_2| = 8 - 6 = 2 \neq 0$

$$A_3 = \begin{bmatrix} 1 & 2 & 4 \\ 3 & 8 & 14 \\ 2 & 6 & 13 \end{bmatrix}$$

$|A_3| = 6 \neq 0.$

→ If the determinant of all the leading submatrices $\neq 0$ then the LU decomposition of that matrix exists otherwise the LU-decomposition of that matrix does not exists.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

e.g

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 5 \\ 1 & 3 & 4 \end{bmatrix}$$

$A_0 = 1$

$|A_0| = 1$

$$A_2 = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \quad |A_2| = 0 \Rightarrow \text{0}.$$

As a hack we can swap R_2 and R_3 then in this case ...

$$A_2 = \begin{bmatrix} 1 & 2 \\ 1 & 3 \end{bmatrix} \quad |A_2| \neq 0.$$

if $|A_3|$ was $=0$ then we cannot apply any hack.

→ LU Decomposition is used as a part of many computations in tools different systems.

→ LU decomposition is not applicable if A is not invertable or $|A_3|=0$.

Gate.Appliedcourse.com

Gatecse@appliedcourse.com

Ph: +91 844-844-0102

Gate.Appliedcourse.com

Gatecse@appliedcourse.com